

ARE FIN WHALES RESIDENT TO THE GULF OF CALIFORNIA?

Bernie R. Tershy¹, Jorge Urbán-Ramírez², Dawn Breese³, Lorenzo Rojas-Bracho⁴ and
Lloyd T. Findley⁵

¹Centro de Conservación y Aprovechamiento de los Recursos Naturales, ITESM-Campus Guaymas, AP 484, Guaymas, Son 85400, México; and Section of Neurobiology and Behavior, Cornell University, Ithaca, NY 14853-2702 USA

²Programa de Mamíferos Marinos, Departamento de Biología Marina, Universidad Autónoma de Baja California Sur, Apdo. Postal 19-B, La Paz, B.C.S., 23000 México

³177 Burns Rd, Brooktondale, NY 14817

⁴Laboratorio Genética, Instituto de Investigaciones Oceanológicas, Universidad Autónoma de Baja California, Apdo. Postal 453, Ensenada, B.C., México.

⁵Departamento de Ciencias Marinas, Instituto Tecnológico y de Estudios Superiores de Monterrey-Campus Guaymas, Apdo. Postal 484, Guaymas, Son, México.

Fin whales (*Balaenoptera physalus*) are relatively abundant in the Gulf of California, México and most technical papers, field guides, and popular articles report that this population may be, or appears to be, resident (e.g. Tomilin 1967, Johnson 1972, Balcomb *et al.* 1979, Wells *et al.* 1981, Leatherwood *et al.* 1982, Gambell 1985).

The earliest reference we could find to a resident Gulf of California fin whale population is Gilmore (1957 p. 23) "...finbacks inhabit the upper Gulf in numbers all year in what appears to be a land locked population". Gilmore's observation, the presence of fin whales in the Gulf of California throughout the year, and the paucity of sightings near the tip of Baja California appear to be the source of subsequent references (S. Leatherwood pers. comm.). In this note we examine existing evidence addressing the hypothesis that fin whales in the Gulf of California comprise a resident population.

Sightings off the Pacific coast of Baja California: According to Rice (1974) the

southern limit of the fin whale's distribution in the Eastern North Pacific is 22°50' N latitude (west of Cabo San Lucas, B.C.S.) during winter, and southern California, sometimes central Baja California, during summer. Urbán and Aguayo (1985) did not record fin whales among 15 cetacean species sighted during 10 cruises along the west coast of Baja California between 1982 and 1985. There were no sightings of fin whales off the Pacific coast of Baja California in the National Marine Fisheries Service data base of cetacean sightings (data provided by S. Reilly). However, the same data base had >25 sightings of blue whales *B. musculus* off the Pacific coast of Baja California Sur (Reilly and Thayer 1990). Blue whales are known to migrate between the Gulf of California and the eastern north Pacific (Tershy *et al.* 1990).

Sightings in the southern Gulf of California: Rojas (1984) reviewed sightings of fin whales in the Gulf from 1957 to 1983 and found none south of 24° N latitude (parallel to La Paz, B.C.S.). Between 1981 and 1987 researchers of the Facultad de Ciencias of the Universidad

Autónoma de México made numerous surveys for cetaceans across the mouth of the Gulf (between Cabo San Lucas, B.C.S. and Mazatlan, Sin. or Cabo Corrientes, Jal.) during all the months of the year and did not sight any fin whales (Aguayo *et al.* 1986, Urbán *et al.* 1988). No fin whales have been sighted during a 5-year study (1988-1992) of humpback whales *Megaptera novaeangliae* at the tip of the Baja California Peninsula (Urbán unpubl. data), however, research was conducted from January-April when one would not expect fin whales to be moving between the Gulf and the Pacific.

Seasonal distribution in the Gulf of California: Fin whales can be found in the Gulf of California in all months of the year. However, 78% of sightings in the southern Gulf (Urbán *et al.*, 1988), 70% of sightings in the Canal de Ballenas, central Gulf (Tershy *et al.*, 1990), 80% of sightings from Isla San Pedro Mártir, central Gulf (Tershy and Breese unpubl. data), 98% of sightings at Guaymas, Sonora, central Gulf (Findley, Vidal and Torre unpubl. data), and 83% of sightings in the northern Gulf (Rojas 1984), were in winter and spring.

Fin whales are more abundant in the Canal de Ballenas area than in other parts of the Gulf (Aguayo *et al.* 1986). In the Canal de Ballenas area summer sea surface temperatures are 2-5° lower (e.g. 25.5° in the Canal de Ballenas area, but 28-29.5° in the rest of the Gulf), and summer primary productivity is higher, than in other parts of the Gulf (Badan-Dangon *et al.* 1985, Alvarez-Borrego 1983). Thus, fin whales in the Gulf during the summer might be expected to concentrate in the Canal de Ballenas area.

Diet and prey availability: Existing evidence suggests that in the Gulf of California fin whales are primarily planktivorous, their most important prey probably being the euphausiid *Nyctiphanes simplex* (Tershy *et al.* in litt.). *N. simplex* is a temperate species and in the

Gulf of California is most abundant in the winter and least abundant in the summer (Brinton *et al.* 1986, Gendron 1990).

Swimming speed and movements in other areas: Southern hemisphere fin whales are thought to migrate 3,800 Km, from breeding grounds to Antarctic feeding grounds, in 1 month, about 130 Km/day (Laws 1961, Gambell 1985). Ray *et al.* (1978) tracked a radio tagged fin whale at an average speed of about 9 Km/hr. Watkins (1981) tracked a radio tagged fin whale which traveled 292 Km in one day, and 2,095 Km in 10 days. Thus, fin whales appear to travel between 100 and 300 Km/day and could thus swim the 1,000 Km length of the Gulf in roughly 3-10 days.

In summary, the available data neither support nor refute the hypothesis that the fin whales in the Gulf of California are a resident or isolated population. Rather, they present a paradox. Sighting data from the southern Gulf and the Pacific coast of Baja California give no indication that fin whales migrate between the Gulf of California and the adjacent Pacific. However, data on the seasonal distribution of fin whales in the Gulf show a marked decline during the summer, when limited evidence suggest that their primary prey is least abundant.

This apparent paradox can be resolved by combining more systematic sighting cruises in the Gulf of California and adjacent Pacific with genetic stock identification studies, comparison of photo-identifications of fin whales from the Pacific with the Gulf of California catalogue (Tershy *et al.* 1990), or satellite tagging of fin whales in the Gulf of California during the late spring. Until these data become available, Gilmore's (1957) hypothesis that fin whales are resident to the Gulf of California should be viewed as untested.

ACKNOWLEDGEMENTS

We thank S. Reilly, NMFS/SWFC, for providing fin whale sightings from the NMFS data base, and S. Leatherwood for insight into the origin of the hypothesis.

LITERATURE CITED

- AGUAYO-L, A., J. URBÁN-R, R. SÁNCHEZ-T, AND L. ROJAS-B. 1986. Diversity and distribution of the cetaceans in the Golfo de California. Abstracts of the XI Reunión Internacional sobre Mamíferos Marinos, 2-6 April, Guaymas, Son., México.
- ALVAREZ-BORREGO, S. 1983. Gulf of California. Pages 427-449 in B. H. Ketchum, ed. Estuaries and enclosed seas. Elsevier Press, Amsterdam.
- BADAN-DANGON, A., C. J. KOBLINSKY, AND T. BAUMGARTNER. 1985. Spring and summer in the Gulf of California: observations of surface thermal patterns. *Oceanologica Acta*, 8:13-22.
- BALCOMB, K.C., B. VILLA-R, AND G. NICHOLS. 1979. Marine mammals in the Sea of Cortez. Proceedings of the 4th international conference on gray whales, La Paz, 16 February 1979.
- BRINTON, E., A. FLEMINGER, AND D. SIEGEL-CAUSEY. 1986. The temperate and tropical planktonic biotas of the Gulf of California. *California Cooperative Investigative Fisheries Reports*, 27:228-266.
- GAMBELL, R. 1985. Fin whale *Balaenoptera physalus* (Linnaeus, 1758). Pages 171-192 in S.H. Ridgway and R. Harrison, eds. The handbook of marine mammals. Volume 3. Academic Press, London.
- GENDRON, D. 1990. Relación entre la abundancia de eufausidos y de ballenas azules (*Balaenoptera musculus*) en el Golfo de California. M.S. thesis, Instituto Politecnico Nacional, Centro Interdisciplinario de Ciencias Marinas, La Paz, BCS, México.
- GILMORE, R.M. 1957. Whales aground in the Cortez Sea. *Pacific Discovery* 10:(1) 22-27.
- JOHNSON, W.W. 1972. Baja California. Time-Life Books, NY
- LAWS, R.M. 1961. Reproduction, growth, and age of southern fin whales. *Discovery Reports* 31:327-486.
- LEATHERWOOD, S., R.R. REEVES, W.F. PERRIN, AND W.E. EVENS. 1982. Whales, dolphins, and porpoises of the eastern north Pacific and adjacent arctic waters. NOAA Technical Report, NMFS Circular 444.
- RAY, G.C., E.D. MITCHEL, D. WARTZOK, V.M. KOZICKI, AND R. MAIEFSKI. 1978. Radio tracking of a fin whale (*Balaenoptera physalus*). *Science* 202:521-524.
- REILLY, S.B. AND V. G. THAYER. 1990. Blue whale (*Balaenoptera musculus*) distribution in the Eastern Tropical Pacific. *Marine Mammal Science* 6:265-277.
- RICE, D.W. 1974. Whales and whale research in the eastern north Pacific. Pages 170-195 in W.E. Schevill, ed. The whale problem: a status report. Harvard University Press, Cambridge.
- ROJAS, B.L. 1984. Presencia y distribución del rorcual común *Balaenoptera physalus* en el Golfo de California, México. Tesis profesional. Facultad de Ciencias, Universidad Autónoma de México.

TERSHEY, B.R., D. BREESE, AND C.S. STRONG. 1990. Abundance, seasonal distribution and population composition of balaenopterid whales in the Canal de Ballenas, Gulf of California, México. Reports of the International Whaling Commission, Special Issue 12:369-375.

TOMLIN, A.G. 1967. Mammals of the USSR and Adjacent Countries, Volume 9 Cetacea. Israel Program for Scientific Translations, Jerusalem.

URBÁN-R, AND A. AGUAYO-L. 1985. Cetáceos observados en la costa occidental de la Península de Baja California, México. Septiembre 1981 - enero 1985. Proceedings of the X Reunión Internacional sobre Mamíferos Marinos 93-118.

URBÁN-R, J., D. AURIOLES-G, AND A. AGUAYO-L. 1988. The fin whale (*Balaenoptera physalus*) in the southern Gulf of California. Abstracts of the XIII Reunión Internacional sobre Mamíferos Marinos, 6-9 April, La Paz, BCS, México.

WATKINS, W.A. 1981. Radio tagging of finback whales- Iceland, June-July 1980. Reference Number 81-82, Woods Hole Oceanographic Institution, Woods Hole, MA.

WELLS, R.S., B.G. WÜRSIG, AND K.S. NORRIS. 1981. A survey of the marine mammals of the upper Gulf of California, Mexico, with an assessment of the status of *Phocoena sinus*. National Technical Information Service Report, PB 81-168791.

Revista de Investigación Científica

PUBLICADA POR LA UNIVERSIDAD AUTONOMA DE BAJA CALIFORNIA SUR
SERIE CIENCIAS DEL MAR